

Search for High-Mass Resonances Decaying to $e\mu$ in pp Collisions at $\sqrt{s}=1.96$ TeV

CDF Collaboration

CAMPANELLI, Mario (Collab.), *et al.*

Abstract

We describe a general search for resonances decaying to a neutral $e\mu$ final state in pp collisions at a center-of-mass energy of 1.96 TeV. Using a data sample representing 344 pb⁻¹ of integrated luminosity recorded by the Collider Detector at Fermilab II experiment, we compare standard model predictions with the number of observed events for invariant masses between 50 and 800 GeV/c². Finding no significant excess (5 events observed vs 7.7 ± 0.8 expected for $M_{e\mu} > 100$ GeV/c²), we set limits on sneutrino and Z' masses as functions of lepton family number violating couplings.

Reference

CDF Collaboration, CAMPANELLI, Mario (Collab.), *et al.* Search for High-Mass Resonances Decaying to $e\mu$ in pp Collisions at $\sqrt{s}=1.96$ TeV. *Physical Review Letters*, 2006, vol. 96, no. 21, p. 211802

DOI : 10.1103/PhysRevLett.96.211802

Available at:

<http://archive-ouverte.unige.ch/unige:38339>

Disclaimer: layout of this document may differ from the published version.



UNIVERSITÉ
DE GENÈVE

Search for High-Mass Resonances Decaying to $e\mu$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV

A. Abulencia,²³ D. Acosta,¹⁷ J. Adelman,¹³ T. Affolder,¹⁰ T. Akimoto,⁵⁵ M. G. Albrow,¹⁶ D. Ambrose,¹⁶ S. Amerio,⁴³ D. Amidei,³⁴ A. Anastassov,⁵² K. Anikeev,¹⁶ A. Annovi,¹⁸ J. Antos,¹ M. Aoki,⁵⁵ G. Apollinari,¹⁶ J.-F. Arguin,³³ T. Arisawa,⁵⁷ A. Artikov,¹⁴ W. Ashmanskas,¹⁶ A. Attal,⁸ F. Azfar,⁴² P. Azzi-Bacchetta,⁴³ P. Azzurri,⁴⁶ N. Bacchetta,⁴³ H. Bachacou,²⁸ W. Badgett,¹⁶ A. Barbaro-Galtieri,²⁸ V. E. Barnes,⁴⁸ B. A. Barnett,²⁴ S. Baroiant,⁷ V. Bartsch,³⁰ G. Bauer,³² F. Bedeschi,⁴⁶ S. Behari,²⁴ S. Belforte,⁵⁴ G. Bellettini,⁴⁶ J. Bellinger,⁵⁹ A. Belloni,³² E. Ben Haim,⁴⁴ D. Benjamin,¹⁵ A. Beretvas,¹⁶ J. Beringer,²⁸ T. Berry,²⁹ A. Bhatti,⁵⁰ M. Binkley,¹⁶ D. Bisello,⁴³ R. E. Blair,² C. Blocker,⁶ B. Blumenfeld,²⁴ A. Bocci,¹⁵ A. Bodek,⁴⁹ V. Boisvert,⁴⁹ G. Bolla,⁴⁸ A. Bolshov,³² D. Bortoletto,⁴⁸ J. Boudreau,⁴⁷ A. Boveia,¹⁰ B. Brau,¹⁰ C. Bromberg,³⁵ E. Brubaker,¹³ J. Budagov,¹⁴ H. S. Budd,⁴⁹ S. Budd,²³ K. Burkett,¹⁶ G. Busetto,⁴³ P. Bussey,²⁰ K. L. Byrum,² S. Cabrera,¹⁵ M. Campanelli,¹⁹ M. Campbell,³⁴ F. Canelli,⁸ A. Canepa,⁴⁸ D. Carlsmith,⁵⁹ R. Carosi,⁴⁶ S. Carron,¹⁵ M. Casarsa,⁵⁴ A. Castro,⁵ P. Catastini,⁴⁶ D. Cauz,⁵⁴ M. Cavalli-Sforza,³ A. Cerri,²⁸ L. Cerrito,⁴² S. H. Chang,²⁷ J. Chapman,³⁴ Y. C. Chen,¹ M. Chertok,⁷ G. Chiarelli,⁴⁶ G. Chlachidze,¹⁴ F. Chlebana,¹⁶ I. Cho,²⁷ K. Cho,²⁷ D. Chokheli,¹⁴ J. P. Chou,²¹ P. H. Chu,²³ S. H. Chuang,⁵⁹ K. Chung,¹² W. H. Chung,⁵⁹ Y. S. Chung,⁴⁹ M. Ciljak,⁴⁶ C. I. Ciobanu,²³ M. A. Ciocci,⁴⁶ A. Clark,¹⁹ D. Clark,⁶ M. Coca,¹⁵ G. Compostella,⁴³ M. E. Convery,⁵⁰ J. Conway,⁷ B. Cooper,³⁰ K. Copic,³⁴ M. Cordelli,¹⁸ G. Cortiana,⁴³ F. Cresciolo,⁴⁶ A. Cruz,¹⁷ C. Cuenca Almenar,⁷ J. Cuevas,¹¹ R. Culbertson,¹⁶ D. Cyr,⁵⁹ S. DaRonco,⁴³ S. D'Auria,²⁰ M. D'Onofrio,³ D. Dagenhart,⁶ P. de Barbaro,⁴⁹ S. De Cecco,⁵¹ A. Deisher,²⁸ G. De Lentdecker,⁴⁹ M. Dell'Orso,⁴⁶ F. Delli Paoli,⁴³ S. Demers,⁴⁹ L. Demortier,⁵⁰ J. Deng,¹⁵ M. Deninno,⁵ D. De Pedis,⁵¹ P. F. Derwent,¹⁶ C. Dionisi,⁵¹ J. R. Dittmann,⁴ P. DiTuro,⁵² C. Dörr,²⁵ S. Donati,⁴⁶ M. Donega,¹⁹ P. Dong,⁸ J. Donini,⁴³ T. Dorigo,⁴³ S. Dube,⁵² K. Ebina,⁵⁷ J. Efron,³⁹ J. Ehlers,¹⁹ R. Erbacher,⁷ D. Errede,²³ S. Errede,²³ R. Eusebi,¹⁶ H. C. Fang,²⁸ S. Farrington,²⁹ I. Fedorko,⁴⁶ W. T. Fedorko,¹³ R. G. Feild,⁶⁰ M. Feindt,²⁵ J. P. Fernandez,³¹ R. Field,¹⁷ G. Flanagan,⁴⁸ L. R. Flores-Castillo,⁴⁷ A. Foland,²¹ S. Forrester,⁷ G. W. Foster,¹⁶ M. Franklin,²¹ J. C. Freeman,²⁸ I. Furic,¹³ M. Gallinaro,⁵⁰ J. Galyardt,¹² J. E. Garcia,⁴⁶ M. Garcia Sciveres,²⁸ A. F. Garfinkel,⁴⁸ C. Gay,⁶⁰ H. Gerberich,²³ D. Gerdes,³⁴ S. Giagu,⁵¹ P. Giannetti,⁴⁶ A. Gibson,²⁸ K. Gibson,¹² C. Ginsburg,¹⁶ N. Giokaris,¹⁴ K. Giolo,⁴⁸ M. Giordani,⁵⁴ P. Giromini,¹⁸ M. Giunta,⁴⁶ G. Giurgiu,¹² V. Glagolev,¹⁴ D. Glenzinski,¹⁶ M. Gold,³⁷ N. Goldschmidt,³⁴ J. Goldstein,⁴² G. Gomez,¹¹ G. Gomez-Ceballos,¹¹ M. Goncharov,⁵³ O. González,³¹ I. Gorelov,³⁷ A. T. Goshaw,¹⁵ Y. Gotra,⁴⁷ K. Goulianos,⁵⁰ A. Gresele,⁴³ M. Griffiths,²⁹ S. Grinstein,²¹ C. Grosso-Pilcher,¹³ R. C. Group,¹⁷ U. Grundler,²³ J. Guimaraes da Costa,²¹ Z. Gunay-Unalan,³⁵ C. Haber,²⁸ S. R. Hahn,¹⁶ K. Hahn,⁴⁵ E. Halkiadakis,⁵² A. Hamilton,³³ B.-Y. Han,⁴⁹ J. Y. Han,⁴⁹ R. Handler,⁵⁹ F. Happacher,¹⁸ K. Hara,⁵⁵ M. Hare,⁵⁶ S. Harper,⁴² R. F. Harr,⁵⁸ R. M. Harris,¹⁶ K. Hatakeyama,⁵⁰ J. Hauser,⁸ C. Hays,¹⁵ A. Heijboer,⁴⁵ B. Heinemann,²⁹ J. Heinrich,⁴⁵ M. Herndon,⁵⁹ D. Hidas,¹⁵ C. S. Hill,¹⁰ D. Hirschbuehl,²⁵ A. Hocker,¹⁶ A. Holloway,²¹ S. Hou,¹ M. Houlden,²⁹ S.-C. Hsu,⁹ B. T. Huffman,⁴² R. E. Hughes,³⁹ J. Huston,³⁵ J. Incandela,¹⁰ G. Introzzi,⁴⁶ M. Iori,⁵¹ Y. Ishizawa,⁵⁵ A. Ivanov,⁷ B. Iyutin,³² E. James,¹⁶ D. Jang,⁵² B. Jayatilaka,³⁴ D. Jeans,⁵¹ H. Jensen,¹⁶ E. J. Jeon,²⁷ S. Jindariani,¹⁷ M. Jones,⁴⁸ K. K. Joo,²⁷ S. Y. Jun,¹² T. R. Junk,²³ T. Kamon,⁵³ J. Kang,³⁴ P. E. Karchin,⁵⁸ Y. Kato,⁴¹ Y. Kemp,²⁵ R. Kephart,¹⁶ U. Kerzel,²⁵ V. Khotilovich,⁵³ B. Kilminster,³⁹ D. H. Kim,²⁷ H. S. Kim,²⁷ J. E. Kim,²⁷ M. J. Kim,¹² S. B. Kim,²⁷ S. H. Kim,⁵⁵ Y. K. Kim,¹³ L. Kirsch,⁶ S. Klimenko,¹⁷ M. Klute,³² B. Knuteson,³² B. R. Ko,¹⁵ H. Kobayashi,⁵⁵ K. Kondo,⁵⁷ D. J. Kong,²⁷ J. Konigsberg,¹⁷ A. Korytov,¹⁷ A. V. Kotwal,¹⁵ A. Kovalev,⁴⁵ A. Kraan,⁴⁵ J. Kraus,²³ I. Kravchenko,³² M. Kreps,²⁵ J. Kroll,⁴⁵ N. Krumnack,⁴ M. Kruse,¹⁵ V. Krutelyov,⁵³ S. E. Kuhlmann,² Y. Kusakabe,⁵⁷ S. Kwang,¹³ A. T. Laasanen,⁴⁸ S. Lai,³³ S. Lami,⁴⁶ S. Lammel,¹⁶ M. Lancaster,³⁰ R. L. Lander,⁷ K. Lannon,³⁹ A. Lath,⁵² G. Latino,⁴⁶ I. Lazzizzera,⁴³ T. LeCompte,² J. Lee,⁴⁹ J. Lee,²⁷ Y. J. Lee,²⁷ S. W. Lee,⁵³ R. Lefèvre,³ N. Leonardo,³² S. Leone,⁴⁶ S. Levy,¹³ J. D. Lewis,¹⁶ C. Lin,⁶⁰ C. S. Lin,¹⁶ M. Lindgren,¹⁶ E. Lipeles,⁹ A. Lister,¹⁹ D. O. Litvintsev,¹⁶ T. Liu,¹⁶ N. S. Lockyer,⁴⁵ A. Loginov,³⁶ M. Loret, ⁴³ P. Loverre,⁵¹ R.-S. Lu,¹ D. Lucchesi,⁴³ P. Lujan,²⁸ P. Lukens,¹⁶ G. Lungu,¹⁷ L. Lyons,⁴² J. Lys,²⁸ R. Lysak,¹ E. Lytken,⁴⁸ P. Mack,²⁵ D. MacQueen,³³ R. Madrak,¹⁶ K. Maeshima,¹⁶ T. Maki,²² P. Maksimovic,²⁴ S. Malde,⁴² G. Manca,²⁹ F. Margaroli,⁵ R. Marginean,¹⁶ C. Marino,²³ A. Martin,⁶⁰ V. Martin,³⁸ M. Martínez,³ T. Maruyama,⁵⁵ H. Matsunaga,⁵⁵ M. E. Mattson,⁵⁸ R. Mazini,³³ P. Mazzanti,⁵ K. S. McFarland,⁴⁹ P. McIntyre,⁵³ R. McNulty,²⁹ A. Mehta,²⁹ S. Menzemer,¹¹ A. Menzione,⁴⁶ P. Merkel,⁴⁸ C. Mesropian,⁵⁰ A. Messina,⁵¹ M. von der Mey,⁸ T. Miao,¹⁶ N. Miladinovic,⁶ J. Miles,³² R. Miller,³⁵ J. S. Miller,³⁴ C. Mills,¹⁰ M. Milnik,²⁵ R. Miquel,²⁸ A. Mitra,¹ G. Mitselmakher,¹⁷ A. Miyamoto,²⁶ N. Moggi,⁵ B. Mohr,⁸ R. Moore,¹⁶ M. Morello,⁴⁶ P. Movilla Fernandez,²⁸ J. Mülmenstädt,²⁸ A. Mukherjee,¹⁶ Th. Muller,²⁵ R. Mumford,²⁴ P. Murat,¹⁶ J. Nachtman,¹⁶ J. Naganoma,⁵⁷ S. Nahn,³² I. Nakano,⁴⁰ A. Napier,⁵⁶ D. Naumov,³⁷ V. Necula,¹⁷ C. Neu,⁴⁵ M. S. Neubauer,⁹ J. Nielsen,²⁸ T. Nigmanov,⁴⁷ L. Nodulman,² O. Norniella,³ E. Nurse,³⁰ T. Ogawa,⁵⁷ S. H. Oh,¹⁵

Y. D. Oh,²⁷ T. Okusawa,⁴¹ R. Oldeman,²⁹ R. Orava,²² K. Osterberg,²² C. Pagliarone,⁴⁶ E. Palencia,¹¹ R. Paoletti,⁴⁶ V. Papadimitriou,¹⁶ A. A. Paramonov,¹³ B. Parks,³⁹ S. Pashapour,³³ J. Patrick,¹⁶ G. Pauletta,⁵⁴ M. Paulini,¹² C. Paus,³² D. E. Pellett,⁷ A. Penzo,⁵⁴ T. J. Phillips,¹⁵ G. Piacentino,⁴⁶ J. Piedra,⁴⁴ L. Pinera,¹⁷ K. Pitts,²³ C. Plager,⁸ L. Pondrom,⁵⁹ X. Portell,³ O. Poukhov,¹⁴ N. Pounder,⁴² F. Prakoshyn,¹⁴ A. Pronko,¹⁶ J. Proudfoot,² F. Ptohos,¹⁸ G. Punzi,⁴⁶ J. Pursley,²⁴ J. Rademacker,⁴² A. Rahaman,⁴⁷ A. Rakitin,³² S. Rappoccio,²¹ F. Ratnikov,⁵² B. Reisert,¹⁶ V. Rekovic,³⁷ N. van Remortel,²² P. Renton,⁴² M. Rescigno,⁵¹ S. Richter,²⁵ F. Rimondi,⁵ L. Ristori,⁴⁶ W. J. Robertson,¹⁵ A. Robson,²⁰ T. Rodrigo,¹¹ E. Rogers,²³ S. Rolli,⁵⁶ R. Roser,¹⁶ M. Rossi,⁵⁴ R. Rossin,¹⁷ C. Rott,⁴⁸ A. Ruiz,¹¹ J. Russ,¹² V. Rusu,¹³ H. Saarikko,²² S. Sabik,³³ A. Safonov,⁵³ W. K. Sakumoto,⁴⁹ G. Salamanna,⁵¹ O. Saltó,³ D. Saltzberg,⁸ C. Sanchez,³ L. Santi,⁵⁴ S. Sarkar,⁵¹ L. Sartori,⁴⁶ K. Sato,⁵⁵ P. Savard,³³ A. Savoy-Navarro,⁴⁴ T. Scheidle,²⁵ P. Schlabach,¹⁶ E. E. Schmidt,¹⁶ M. P. Schmidt,⁶⁰ M. Schmitt,³⁸ T. Schwarz,³⁴ L. Scodellaro,¹¹ A. L. Scott,¹⁰ A. Scribano,⁴⁶ F. Scuri,⁴⁶ A. Sedov,⁴⁸ S. Seidel,³⁷ Y. Seiya,⁴¹ A. Semenov,¹⁴ L. Sexton-Kennedy,¹⁶ I. Sfiligoi,¹⁸ M. D. Shapiro,²⁸ T. Shears,²⁹ P. F. Shepard,⁴⁷ D. Sherman,²¹ M. Shimojima,⁵⁵ M. Shochet,¹³ Y. Shon,⁵⁹ I. Shreyber,³⁶ A. Sidoti,⁴⁴ P. Sinervo,³³ A. Sisakyan,¹⁴ J. Sjolin,⁴² A. Skiba,²⁵ A. J. Slaughter,¹⁶ K. Sliwa,⁵⁶ J. R. Smith,⁷ F. D. Snider,¹⁶ R. Snihur,³³ M. Soderberg,³⁴ A. Soha,⁷ S. Somalwar,⁵² V. Sorin,³⁵ J. Spalding,¹⁶ M. Spezziga,¹⁶ F. Spinella,⁴⁶ T. Spreitzer,³³ P. Squillacioti,⁴⁶ M. Stanitzki,⁶⁰ A. Staveris-Polykalas,⁴⁶ R. St. Denis,²⁰ B. Stelzer,⁸ O. Stelzer-Chilton,⁴² D. Stentz,³⁸ J. Strologas,³⁷ D. Stuart,¹⁰ J. S. Suh,²⁷ A. Sukhanov,¹⁷ K. Sumorok,³² H. Sun,⁵⁶ T. Suzuki,⁵⁵ A. Taffard,²³ R. Takashima,⁴⁰ Y. Takeuchi,⁵⁵ K. Takikawa,⁵⁵ M. Tanaka,² R. Tanaka,⁴⁰ N. Tanimoto,⁴⁰ M. Tecchio,³⁴ P. K. Teng,¹ K. Terashi,⁵⁰ S. Tether,³² J. Thom,¹⁶ A. S. Thompson,²⁰ E. Thomson,⁴⁵ P. Tipton,⁴⁹ V. Tiwari,¹² S. Tkaczyk,¹⁶ D. Toback,⁵³ S. Tokar,¹⁴ K. Tollefson,³⁵ T. Tomura,⁵⁵ D. Tonelli,⁴⁶ M. Tönnemann,³⁵ S. Torre,¹⁸ D. Torretta,¹⁶ S. Tourneur,⁴⁴ W. Trischuk,³³ R. Tsuchiya,⁵⁷ S. Tsuno,⁴⁰ N. Turini,⁴⁶ F. Ukegawa,⁵⁵ T. Unverhau,²⁰ S. Uozumi,⁵⁵ D. Usynin,⁴⁵ A. Vaiciulis,⁴⁹ S. Vallecorsa,¹⁹ A. Varganov,³⁴ E. Vataga,³⁷ G. Velev,¹⁶ G. Veramendi,²³ V. Veszpremi,⁴⁸ R. Vidal,¹⁶ I. Vila,¹¹ R. Vilar,¹¹ T. Vine,³⁰ I. Vollrath,³³ I. Volobouev,²⁸ G. Volpi,⁴⁶ F. Würthwein,⁹ P. Wagner,⁵³ R. G. Wagner,² R. L. Wagner,¹⁶ W. Wagner,²⁵ R. Wallny,⁸ T. Walter,²⁵ Z. Wan,⁵² S. M. Wang,¹ A. Warburton,³³ S. Waschke,²⁰ D. Waters,³⁰ W. C. Wester III,¹⁶ B. Whitehouse,⁵⁶ D. Whiteson,⁴⁵ A. B. Wicklund,² E. Wicklund,¹⁶ G. Williams,³³ H. H. Williams,⁴⁵ P. Wilson,¹⁶ B. L. Winer,³⁹ P. Wittich,¹⁶ S. Wolbers,¹⁶ C. Wolfe,¹³ T. Wright,³⁴ X. Wu,¹⁹ S. M. Wynne,²⁹ A. Yagil,¹⁶ K. Yamamoto,⁴¹ J. Yamaoka,⁵² T. Yamashita,⁴⁰ C. Yang,⁶⁰ U. K. Yang,¹³ Y. C. Yang,²⁷ W. M. Yao,²⁸ G. P. Yeh,¹⁶ J. Yoh,¹⁶ K. Yorita,¹³ T. Yoshida,⁴¹ G. B. Yu,⁴⁹ I. Yu,²⁷ S. S. Yu,¹⁶ J. C. Yun,¹⁶ L. Zanello,⁵¹ A. Zanetti,⁵⁴ I. Zaw,²¹ F. Zetti,⁴⁶ X. Zhang,²³ J. Zhou,⁵² and S. Zucchelli⁵

(CDF Collaboration)

¹*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*²*Argonne National Laboratory, Argonne, Illinois 60439, USA*³*Institut de Fisica d'Altes Energies, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*⁴*Baylor University, Waco, Texas 76798, USA*⁵*Istituto Nazionale di Fisica Nucleare, University of Bologna, I-40127 Bologna, Italy*⁶*Brandeis University, Waltham, Massachusetts 02254, USA*⁷*University of California, Davis, Davis, California 95616, USA*⁸*University of California, Los Angeles, Los Angeles, California 90024, USA*⁹*University of California, San Diego, La Jolla, California 92093, USA*¹⁰*University of California, Santa Barbara, Santa Barbara, California 93106, USA*¹¹*Instituto de Fisica de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*¹²*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*¹³*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*¹⁴*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*¹⁵*Duke University, Durham, North Carolina 27708*¹⁶*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*¹⁷*University of Florida, Gainesville, Florida 32611, USA*¹⁸*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*¹⁹*University of Geneva, CH-1211 Geneva 4, Switzerland*²⁰*Glasgow University, Glasgow G12 8QQ, United Kingdom*²¹*Harvard University, Cambridge, Massachusetts 02138, USA*²²*Division of High Energy Physics, Department of Physics, University of Helsinki, Helsinki, Finland and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland*²³*University of Illinois, Urbana, Illinois 61801, USA*

- ²⁴The Johns Hopkins University, Baltimore, Maryland 21218, USA
²⁵Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany
²⁶High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305, Japan
²⁷Center for High Energy Physics: Kyungpook National University, Taegu 702-701 Korea;
 Seoul National University, Seoul 151-742, Korea;
 and SungKyunKwan University, Suwon 440-746; Korea
²⁸Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA
²⁹University of Liverpool, Liverpool L69 7ZE, United Kingdom
³⁰University College London, London WC1E 6BT, United Kingdom
³¹Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain
³²Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA
³³Institute of Particle Physics: McGill University, Montréal, Canada H3A 2T8;
 and University of Toronto, Toronto, Canada M5S 1A7
³⁴University of Michigan, Ann Arbor, Michigan 48109, USA
³⁵Michigan State University, East Lansing, Michigan 48824, USA
³⁶Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia
³⁷University of New Mexico, Albuquerque, New Mexico 87131, USA
³⁸Northwestern University, Evanston, Illinois 60208, USA
³⁹The Ohio State University, Columbus, Ohio 43210, USA
⁴⁰Okayama University, Okayama 700-8530, Japan
⁴¹Osaka City University, Osaka 588, Japan
⁴²University of Oxford, Oxford OX1 3RH, United Kingdom
⁴³University of Padova, Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, I-35131 Padova, Italy
⁴⁴LPNHE, Universite Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France
⁴⁵University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
⁴⁶Istituto Nazionale di Fisica Nucleare Pisa, Universities of Pisa, Siena and Scuola Normale Superiore, I-56127 Pisa, Italy
⁴⁷University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA
⁴⁸Purdue University, West Lafayette, Indiana 47907, USA
⁴⁹University of Rochester, Rochester, New York 14627, USA
⁵⁰The Rockefeller University, New York, New York 10021, USA
⁵¹Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, University of Rome "La Sapienza," I-00185 Roma, Italy
⁵²Rutgers University, Piscataway, New Jersey 08855, USA
⁵³Texas A&M University, College Station, Texas 77843, USA
⁵⁴Istituto Nazionale di Fisica Nucleare, University of Trieste/Udine, Italy
⁵⁵University of Tsukuba, Tsukuba, Ibaraki 305, Japan
⁵⁶Tufts University, Medford, Massachusetts 02155, USA
⁵⁷Waseda University, Tokyo 169, Japan
⁵⁸Wayne State University, Detroit, Michigan 48201, USA
⁵⁹University of Wisconsin, Madison, Wisconsin 53706, USA
⁶⁰Yale University, New Haven, Connecticut 06520, USA
 (Received 7 March 2006; published 30 May 2006)

We describe a general search for resonances decaying to a neutral $e\mu$ final state in $p\bar{p}$ collisions at a center-of-mass energy of 1.96 TeV. Using a data sample representing 344 pb^{-1} of integrated luminosity recorded by the Collider Detector at Fermilab II experiment, we compare standard model predictions with the number of observed events for invariant masses between 50 and 800 GeV/c^2 . Finding no significant excess (5 events observed vs 7.7 ± 0.8 expected for $M_{e\mu} > 100 \text{ GeV}/c^2$), we set limits on sneutrino and Z' masses as functions of lepton family number violating couplings.

DOI: [10.1103/PhysRevLett.96.211802](https://doi.org/10.1103/PhysRevLett.96.211802)

PACS numbers: 13.85.Rm, 12.60.Cn, 12.60.Jv, 14.70.Pw

An observed excess of high-mass opposite sign electron-muon pairs at the Tevatron would provide evidence of physics beyond the standard model (SM). Nearly every extension of the SM predicts flavor changing neutral current effects. General supersymmetric (SUSY) models, for instance, permit the violation of R -parity symmetry (RPV) and describe the lepton family number violating (LFV) production and decay of sneutrinos ($\tilde{\nu}$), the scalar superpartners of SM neutrinos [1]. Models with additional gauge symmetry can also accommodate an $e\mu$ signature through

LFV decays of a new heavy neutral gauge boson, the Z' [2,3]. Signals from such processes may be easily detected at the Tevatron. SM backgrounds are small and characterized by invariant mass spectra that lie well below the range presumed for new physics.

We search for a general signature of high-mass opposite sign $e\mu$ pairs and are sensitive to both resonant and non-resonant production mechanisms. The CDF collaboration investigated the high-mass $e\mu$ channel in a Run I search for the direct RPV production and decay of the tau sneu-

trino ($\tilde{\nu}_\tau$) [4,5]. That analysis excluded sneutrino masses below $\sim 360 \text{ GeV}/c^2$ for a particular set of RPV coupling values. This Letter describes our continued investigation of this channel using data from Run II of the Tevatron taken with the upgraded Collider Detector at Fermilab (CDF II). The data sample we use represents an integrated luminosity of $344 \pm 21 \text{ pb}^{-1}$, approximately 3 times that used in Run I. We search for an $e\mu$ resonance by examining invariant masses ($M_{e\mu}$) between 100 and $800 \text{ GeV}/c^2$ for an excess of events. We find event counts that are consistent with SM predictions and use our result to constrain models of LFV $\tilde{\nu}$ and Z' decay by excluding $e\mu$ coupling values as functions of the new particle masses. In contrast to searches that utilize low-energy data, where the effects of new heavy particles may be masked by cancellations of various contributions, the interpretation of our results is robust.

The CDF II detector is an azimuthally and forward-backward symmetric apparatus designed to study $p\bar{p}$ reactions at the Tevatron. The detector has a charged particle tracking system immersed in a 1.4 T magnetic field aligned coaxially with the $p\bar{p}$ beams. A 3.1 m long open-cell drift chamber, the Central Outer Tracker [6], covers the radial range from 40 to 137 cm and provides coverage for the pseudorapidity range $|\eta| \lesssim 1$ [7]. Segmented electromagnetic and hadronic sampling calorimeters surround the tracking system and measure the energies of interacting particles in the range $|\eta| < 3.6$ [8–10]. A set of drift chambers located outside the central hadron calorimeters and another set behind a 60 cm thick iron shield track muons with $|\eta| \leq 0.6$ [11]. Additional drift chambers and scintillation counters detect muons in the region $0.6 \leq |\eta| \leq 1.0$. Gas Čerenkov counters located in the $3.7 < |\eta| < 4.7$ region [12] measure the average number of inelastic $p\bar{p}$ collisions per bunch crossing and thereby determine the beam luminosity.

We use data collected with high- P_T triggers that require central ($|\eta| \lesssim 1.0$) lepton candidates. We select events that contain at least one reconstructed electron of $E_T > 20 \text{ GeV}$ and one reconstructed muon with $P_T > 20 \text{ GeV}/c$. Energy deposited by the leptons in the central electromagnetic calorimeters must be of an isolated nature and should also satisfy the standard set of “tight” CDF lepton identification (ID) criteria [13].

We select oppositely charged $e\mu$ pairs and ensure that the lepton tracks come from a common $p\bar{p}$ interaction. We reject events containing cosmic-ray muons and photon-conversion electrons by imposing additional event topology requirements [13]. In order to maximize acceptance to a broad range of new physics signatures, we do not impose requirements on missing transverse energy or jets.

SM $e\mu$ backgrounds include $q\bar{q} \rightarrow Z/\gamma^* \rightarrow \tau\tau$ (Drell-Yan), top ($t\bar{t}$) and diboson (WW , WZ , and ZZ) production. We determine geometric and kinematic acceptances for these processes using the PYTHIA Monte Carlo generator

[14] and a GEANT3 [15] based simulation of the CDF II detector. These simulations employ the CTEQ5L [16] parton distribution functions (PDF’s) to model the momentum distribution of the initial-state partons. We define the acceptance for each background process, α_i , as the fraction of generated events that satisfy the lepton ID and event topology requirements. We correct the α_i by multiplying with trigger efficiencies (ϵ_{trg}) measured from $W \rightarrow e\nu$ and $Z \rightarrow \mu\mu$ data and factors (f_{reco} , f_{ID}) that account for differences in lepton reconstruction and ID efficiency between simulation and data [13]. We refer to the combined correction factor, $\epsilon_{\text{trg}} \times f_{\text{reco}} \times f_{\text{ID}}$, as ϵ .

The expected contribution of each SM background is given by the product of $\alpha_i \times \epsilon$ with the corresponding cross section and the integrated luminosity of our data sample. We estimate top and diboson background contributions using next-to-leading order (NLO) cross sections: 6.1 pb for $t\bar{t}$ [17], 12.1 pb for WW , 3.7 pb for WZ , and 1.4 pb for ZZ [18]. We use a next-to-next-to-leading order continuum ($M_{ll} > 30 \text{ GeV}/c^2$) cross section for the Drell-Yan process, 337.7 pb, which we calculate by scaling the PYTHIA leading order cross section by the ratio of NNLO to LO predictions obtained from PHOZPR calculations [19].

Jets that are misidentified as leptons account for approximately 5% of the total background. We determine the misidentification probability (“fake rate”) [13] by examining separate data samples collected with various jet E_T triggers for the occurrence of leptons. The real leptonic content of these samples is assumed to be negligible. We obtain fake rates from the ratios of misidentified leptons and the number of jets in these samples. We then estimate the background from this source by applying the measured fake rates, parametrized as functions of E_T , to the jets in events in the high- P_T sample that contain a single lepton candidate.

We present the number of observed and predicted background events for two $M_{e\mu}$ regions in Table I. The dominant uncertainty on the background predictions arises from a 6% uncertainty in the luminosity measurement [20]. Additional uncertainty contributions include those associated with the SM cross sections, fake rate measurements, lepton ID and reconstruction scale factors, and PDF model.

TABLE I. Expected and observed event totals for low- and high-mass $M_{e\mu}$ regions. The uncertainties shown are the combination of statistical and systematic errors.

Channel	$50 < M_{e\mu} < 100 \text{ GeV}/c^2$	$M_{e\mu} > 100 \text{ GeV}/c^2$
$Z/\gamma^* \rightarrow \tau\tau$	38.8 ± 2.9	0.6 ± 0.0
WW, WZ, ZZ	6.6 ± 0.5	3.5 ± 0.2
$t\bar{t}$	3.6 ± 0.5	3.2 ± 0.5
Fake Lepton	2.9 ± 1.7	$0.4^{+0.6}_{-0.4}$
Prediction	51.9 ± 3.4	7.7 ± 0.8
Observation	56	5

Overall, we find that uncertainties on our signal acceptance and background expectations have little impact on the limits we set.

The $50 \leq M_{e\mu} \leq 100 \text{ GeV}/c^2$ region listed in Table I represents an invariant mass range that is rich in background. Finding good agreement between our observation and predicted background in this region, we next consider the $M_{e\mu}$ range above $100 \text{ GeV}/c^2$. Here, too, we find good agreement. Figure 1 shows the observed and predicted background $M_{e\mu}$ distributions over a portion of the full $50\text{--}800 \text{ GeV}/c^2$ invariant mass range.

We quantify the consistency between the $M_{e\mu}$ distributions presented in Fig. 1 by performing a χ^2 test, using variable-width $M_{e\mu}$ bins to achieve sufficient predicted background occupancies for the Gaussian approximation of Poisson statistics. The test results in a total reduced χ^2 statistic (χ^2/N_{dof}) of $14.0/11 = 1.27$, under the assumption that systematic uncertainties in each bin are completely correlated. The statistic implies a p value, the probability of finding a larger χ^2 , of $\sim 23\%$. This value provides a statistical basis for accepting our results as consistent with SM expectations.

Finding no evidence of new physics in the $M_{e\mu}$ spectrum, we turn from the model-independent search to constraints on specific models; RPV sneutrino and LFV Z' decay. In RPV SUSY models, the s -channel $d\bar{d} \rightarrow \tilde{\nu}_i(\tilde{\nu}_i) \rightarrow e\mu$ process is governed by two LFV couplings. λ'_{i11} determines the sneutrino production cross section from d and \bar{d} while λ_{i2} gives the sneutrino branching ratio to $e\mu$ [21]. The index i refers to the lepton generation of the sneutrino. We do not consider initial states that include up-type quarks since RPV sneutrino hadro-production occurs only through a $L_i Q_j \bar{D}_k$ term in the superpotential [21]. The final state for this process consists of $e\mu$ only, i.e., it does not contain additional neutrinos or a SUSY lightest supersymmetric particle.




FIG. 1 (color online). Observed and predicted $M_{e\mu}$ Distributions. The observed $e\mu$ invariant mass spectrum agrees well with that of the combined SM and fake lepton backgrounds. No events are observed in data beyond $159 \text{ GeV}/c^2$.

Since strong limits exist for $\tilde{\nu}_e$ and $\tilde{\nu}_\mu$ couplings [22], we focus instead on the third generation sneutrino, $\tilde{\nu}_\tau$. We assume the “single coupling dominance” hypothesis [22] and set all λ and λ' couplings but λ'_{311} and λ_{132} to zero so that contributions to the $e\mu$ channel originate from the $\tilde{\nu}_\tau$ only. Previous limits on λ'_{311} and λ_{132} from low-energy experiments are 0.10 and 0.05 [23], assuming squark and slepton masses of $100 \text{ GeV}/c^2$.

We show an example NLO $\sigma \times \text{BR}$ [24] that corresponds to $\lambda'_{311} = 0.10$ and $\lambda_{132} = 0.05$ in Fig. 2. We use such curves for different λ_{132} and λ'_{311} values and a mass-dependent $\alpha \times \epsilon$ calculated from PYTHIA Monte Carlo simulations of the $d\bar{d} \rightarrow \tilde{\nu}_\tau(\tilde{\nu}_\tau) \rightarrow e\mu$ process to obtain signal expectations over the 100 to $800 \text{ GeV}/c^2$ range. Assuming coupling values of $\lambda_{132} = 0.05$ and $\lambda'_{311} = 0.1$, for example, we find that a hypothetical $300 \text{ GeV}/c^2$ $\tilde{\nu}_\tau$ signal consists of 16.2 ± 1.3 events.

We calculate an upper limit $\sigma \times \text{BR}$ for $d\bar{d} \rightarrow \tilde{\nu}_\tau(\tilde{\nu}_\tau) \rightarrow e\mu$ using the numbers of observed and predicted background events in bins of $M_{e\mu}$. We scan the $M_{e\mu}$ range in steps of $10 \text{ GeV}/c^2$ and use the simulated mass resolution at each step to weight all data and background events between 100 and $800 \text{ GeV}/c^2$. A Bayesian algorithm [25] is used with a uniform prior to translate the weighted event totals to a 95% confidence level (C.L.) upper limit on the number of signal events in data. We divide this limit by the integrated luminosity and signal $\alpha \times \epsilon$ to obtain a 95% C.L. upper limit on the $\sigma \times \text{BR}$ for the process. The $\alpha \times \epsilon$ for a generic spin-0 particle decaying to $e\mu$ is mass dependent, increasing slowly from $\sim 10\%$ at




FIG. 2 (color online). Observed 95% C.L. upper limit on $\sigma \times \text{BR}$ for $d\bar{d} \rightarrow \tilde{\nu}_\tau(\tilde{\nu}_\tau) \rightarrow e\mu$ (solid line) and the NLO prediction (dashed line) as a function of $e\mu$ invariant mass. Their intersection gives a $530 \text{ GeV}/c^2$ $\tilde{\nu}_\tau$ mass limit for the values of λ'_{311} and λ_{132} indicated. Because of small differences in the signal acceptances, the 95% C.L. upper limit on the Z' $\sigma \times \text{BR}$ (not shown) is larger than that of the sneutrino, $\sim 0.02 \text{ pb}$ greater at $200 \text{ GeV}/c^2$ and $\sim 0.003 \text{ pb}$ greater at $600 \text{ GeV}/c^2$, for example.




FIG. 3 (color online). $\lambda'_{311} - M_{e\mu}$ exclusion regions. Regions to the left of the five curves shown represent ranges of λ'_{311} values that we exclude at 95% C.L. as a function of $\tilde{\nu}_\tau$ mass. Each region corresponds to a fixed value of λ_{132} .

100 GeV/c^2 to $\sim 27\%$ at 800 GeV/c^2 . We account for uncertainties on the signal and background expectations in limit calculation although the final results are largely insensitive to these numbers.

The intersection of the observed upper limit and a NLO $\sigma \times \text{BR}$ curve defines a $\tilde{\nu}_\tau$ mass limit for specific values of the λ'_{311} and λ_{132} couplings, as shown in Fig. 2. We construct exclusion regions in the $\lambda'_{311} - M_{e\mu}$ plane by plotting the mass limit as a function of both RPV couplings. Figure 3 shows the exclusion region parametrized by five assumed values of λ_{132} . The plot indicates, for example, that we exclude at 95% C.L. λ'_{311} values above ~ 0.01 for a sample $\tilde{\nu}_\tau$ mass of 300 GeV/c^2 and $\lambda_{132} \geq 0.02$.

Our search is also sensitive to LFV Z' decay. The $p\bar{p} \rightarrow Z' \rightarrow e\mu$ process proceeds through diagonal $U(1)'$ couplings at the initial-state vertex and an off-diagonal LFV $U(1)'$ coupling, Q'_{12} , that determines the Z' branching ratio (BR) to $e\mu$ [2]. We set 95% C.L. limits on Q'_{12} as a function of the Z' mass using NLO $\sigma \times \text{BR}$'s from a group of E_6 -inspired models. Although E_6 models do not incorporate the LFV Q'_{12} coupling by construction, we use these models because they provide a theoretical reference by specifying initial-state Z' -quark coupling and a NLO Z' production cross section. We utilize NLO cross sections provided by the χ , ψ , secluded, and η E_6 models [26] and extend these models to include the Q'_{12} coupling [27], which introduces $Z' \rightarrow e\mu$ decays.

We set limits on Q'_{12} as a function of the Z' mass following the procedure previously described for RPV $\tilde{\nu}_\tau$ decay. The $\alpha \times \epsilon$ we use in calculating the upper limit on $\sigma \times \text{BR}$ is obtained from PYTHIA Monte Carlo simulations in which the Z' is required to couple to a left-handed leptonic LFV current, as may be favored in E_6 -motivated $U(1)'$ models that incorporate LFV [27]. Z' acceptance ($\sim 8\%$ at 100 GeV/c^2 , and increasing to $\sim 20\%$ at




FIG. 4 (color online). $Q'_{12} - M_{e\mu}$ exclusion regions depicting 95% C.L. upper limits on the LFV Q'_{12} coupling in extended ψ , χ , η , and secluded E_6 models.

800 GeV/c^2) is smaller than that for the scalar sneutrino due to the spin-1 nature of the particle.

The $Q'_{12} - M_{e\mu}$ regions that we exclude for the modified χ , ψ , secluded, and η models are shown in Fig. 4. Assuming initial-state Z' couplings specified by the η model, for example, we exclude Q'_{12} values above ~ 0.01 – 0.1 for Z' masses between 200 and 700 GeV/c^2 . Because the LFV Q'_{12} coupling is not intrinsic to E_6 models, our limits should not be interpreted as constraints on the models themselves.

In summary, we searched in 344 pb^{-1} of CDF II data for high-mass $e\mu$ events and found an $M_{e\mu}$ distribution consistent with SM predictions. With no evidence for new physics, we set correlated 95% C.L. limits on the mass and λ'_{311} and λ_{132} RPV couplings of the $\tilde{\nu}_\tau$. We achieve a ~ 170 GeV/c^2 improvement in the $\tilde{\nu}_\tau$ mass limit for specific RPV coupling values of $\lambda'_{311} = 0.1$ and $\lambda_{132} = 0.05$ [28] and, more generally, exclude $\tilde{\nu}_\tau$ masses over a range of λ'_{311} and λ_{132} values. We also place limits on potential LFV Q'_{12} couplings of the Z' boson as a function of its mass using E_6 -like models of $U(1)'$ symmetry.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. We thank S. Mahji for providing us with NLO $\tilde{\nu}$ production cross sections and B. Dobrescu, J. Kang and P. Langacker for guidance on incorporating LFV Z' decay in E_6 models. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium fuer Bildung und Forschung, Germany; the Korean Science and Engineering Foundation and the Korean Research

Foundation; the Particle Physics and Astronomy Research Council and the Royal Society, UK; the Russian Foundation for Basic Research; the Comision Interministerial de Ciencia y Tecnologia, Spain; in part by the European Community's Human Potential Programme under Contract No. HPRN-CT-20002, Probe for New Physics; by the Research Fund of Istanbul University Project No. 1755/21122001; and by the Research Corporation.

-
- [1] S. Yan-Bin *et al.*, Commun. Theor. Phys. **44**, 107 (2005).
 - [2] See, for example, B. Murakami, Phys. Rev. D **65**, 055003 (2002).
 - [3] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. Lett. **96**, 211801 (2006). This paper describes a Z' search that uses the angular distribution of dielectrons.
 - [4] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. Lett. **91**, 171602 (2003).
 - [5] B. Abbott *et al.* (D0 Collaboration), Phys. Rev. D **62**, 092004 (2000). This paper describes a Run I search that includes the $e\mu$ channel.
 - [6] T. Affolder *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **526**, 249 (2004).
 - [7] We use a cylindrical coordinate system about the beam axis in which θ is the polar angle, ϕ is the azimuthal angle, and η is pseudorapidity, $-\ln \tan(\theta/2)$. $E_T \equiv E \sin \theta$ and $P_T \equiv P \sin \theta$ where E is energy measured by the calorimeter and P is the magnitude of the momentum measured by the spectrometer. $\cancel{E}_T \equiv -\sum_i E_T^i \mathbf{n}_i$, where \mathbf{n}_i is the unit vector in the azimuthal plane that points from the beamline to the i th calorimeter tower.
 - [8] L. Balka *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **267**, 272 (1988).
 - [9] S. Bertolucci *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **267**, 301 (1988).
 - [10] M. Albrow *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **480**, 524 (2002).
 - [11] G. Ascoli *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **268**, 33 (1988).
 - [12] D. Acosta *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **461**, 540 (2001).
 - [13] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. D (to be published).
 - [14] Computer code PYTHIA v6_216, in T. Sjöstrand *et al.*, Comput. Phys. Commun. **135**, 238 (2001).
 - [15] Computer code GEANT v3.2113, in R. Brun *et al.*, Report No. CERN-DD-78-2-REV.
 - [16] H.L. Lai *et al.*, Eur. Phys. J. C **12**, 375 (2000).
 - [17] M. Cacciari, S. Frixione, M.L. Mangano, P. Nason, and G. Ridolfi, J. High Energy Phys. 04 (2004) 068.
 - [18] J.M. Campbell and R.K. Ellis, Phys. Rev. D **60**, 113006 (1999).
 - [19] C.R. Hamberg, W.L. van Neerven, and T. Matsuura, Nucl. Phys. **B359**, 343 (1991).
 - [20] S. Klimenko *et al.*, Report No. FERMILAB-FN-0741 (2003); D. Acosta *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **494**, 57 (2002).
 - [21] D. Choudhury, S. Majhi, and V. Ravindran, Nucl. Phys. **B660**, 343 (2003).
 - [22] M. Chemtob, Prog. Part. Nucl. Phys. **54**, 71 (2005).
 - [23] F. Ledroit and G. Sajot, Report No. GDR-S-008 (1998).
 - [24] S. Majhi (private communications).
 - [25] J. Heinrich *et al.* (CDF Statistics Committee), physics/0409129.
 - [26] P. Langacker and J. Kang, Phys. Rev. D **71**, 035014 (2005).
 - [27] P. Langacker (private communications).
 - [28] S. Eidelman *et al.* (Particle Data Group), Phys. Lett. B **592**, 1 (2004).